95 PV 860058 (Ancience Edihan Voin 88/064 Vol 4. Mo. 3 te 86/095



P342

GUE

MIRCEN Journal, 198K, 4, 000-000

and a state of the

to additioned

# Nitrogen fixation in bambara groundnut, Voandzeia subterranea (L.) Thouars CN0-10-1182

Mamadou Gueye<sup>1</sup> and Lucien M. Bordeleau<sup>2</sup>

Received 19 February 1988, accepted 21 March 1988

# Introduction

Bambara groundnut (Voandzeia subterranea (L.) Thouars) is a desirable legume for Africa semi-arid tropical agriculture. Like cowpeas, the bambara groundnut nodulates freely in tropical soils. Although it produces a nutritious food, the bambara groundnut remains one of the crops most neglected by science (National Academy of Sciences, 1979).

The most important feature of legumes is their ability to fix atmospheric nitrogen and considerable efforts have been expended towards increasing the efficiency of its symbiotic association with Rhizobium. Now, it is accepted that seed inoculation with superior *Rhizobium* strains is necessary with most legumes. However, cowpea seed inoculation is usually unsuccessful because of competition between introduced and indigenous *Rhizobium* strains in the soil. Therefore, an important step towards improved nitrogen fixation is to isolate and identify competitive indigenous *Rhizobia* with a view to using elite strains as inoculants (Mulongoy et al., 1980).

This paper presents data on screening field bambara groundnut varieties, nitrogen fixing ability of indigenous Rhizobium strains and evaluation of interactions between host plant and Rhizobium strain.

# Materials and methods

Field varietal screening. Trials were done at two sites in Senegal during the 1985 rainy season. The sites differed in both rainfall (Bambey, 377 mm annum<sup>-1</sup>; Nioro, 532 mm annum<sup>-1</sup>) and in soil type (Table 1). Twenty four cultivars of bambara groundnut (listed'in Fig. 1) were hand sown in a randomized block design replicated four times. The **cultivars** were sown in two treatments: without (-N) and with (+N) nitrogen fertilization (50 kg urea/ha). All plots received 60 kg P<sub>2</sub>O<sub>4</sub>/ha and 120 kg KCI/ha. Six weeks after planting, the relative effectiveness (RE) of each host plant/ indigenous Rhizobium strain combination was evaluated as follows (Mulongoy et al., 1980):

<sup>1</sup>MIRCEN-CNRA, B.P. 53, **Bambey, Senegal.** <sup>2</sup>Canada Agriculture, Research Statron, 2560 Hochelaga **blvd**, Sainte-Foy, Canada, G1V 2J3.

© Oxford University Press 1988

Soil	_	ppm	Composition (%)						
	Total C	Total N	Available P (Olsen*)	Clay	Loam	Sand	pH (H₂O)		
Bambey	2970	290	121	4	2	96	7.x		
Nioro	4160	310	137	6	3	90	5.8		

Table I Characteristics of soil at Bambey and Nioro experimental stations

\*Olsen et al. (1954).

$$RE = \frac{\text{`weight' of ten samples in the -N block}}{\text{`weight' of ten samples in the -t-N block}} \times 100$$

In this formula, 'weight' is either of shoot, **root** or nodule dry weight or shoot **nitrogen** content.

Nitrogen fixing ability of indigenous Rhizobium strains. The samples were taken in each treatment from five well-developed plants and still having immature pods and 45 well-developed, pink nodules were sampled from the tap roots. Rhizobium strains were isolated according to standard procedures (Vincent, 1970). Isolated rhizobia were cultured on agar slants made from yeast extractinaltose medium. Nitrogen fixing ability of the isolates were compared in terms obfsshout dry- weight of bambara groundnut cultivar 85001 used as test plant and cultivated in Leonard jar assemblies (Leonard, 1943). Seeds were sterilized using 0.01% (WV) HgCl<sub>2</sub> for 3 min and germinated in sterile sand for 3 d; each seedling was then transplanted into a jar assembly. One ml suspension of rhizobial isolate broth (approx. 10<sup>9</sup> cells) was added aseptically to the seedlings for ail treatments except for an uninoculated control and plants were grown in a greenhouse. Nitrogen-free Hewitt solution (Hewitt, 1946) diluted with water (1:3 v/v) was added once a week during the growing period. After 35 d growth, plant tops were clipped off, dried and weighed for dry matter production.

**Greenhouse evaluation of host plant-Rhizobium** strain interactions. Both the indigenous and Niftal Rhizobium strains (Iisted in Table 3) were selected on the basis of their effectiveness. The Niftal strains – TAL 22, TAL 169 and TAL 569 – were supplied by the NiftAL project. Hawaii, USA and served as standards for comparison against the indigenous ones. Twelve bambara groundnut cultivars also listed in Table 3 were used as host plants. Seed sterilization of each genotype, germination, transplantation, inoculation and watering were conducted as described above and seeds were planted in Leonard jars. Treatments were arranged in a randomized block design replicated five times. After 50 days of growth, the plants were harvested. Plant tops were dried and weighed. Nodulation was scored taking into account the number (N), the internal colour (C) and the size (S) of nodules. The nodulation index (Nod. I.) was calculated as follows:

#### Nod. I. = $N \times C \times S$

Nodule number was rated on a scale from 0 (no nodule) to 3 (many nodules). Nodule internal colour was from 0 (white) to 1 (red) and nodule size was from 1 (small nodule) to 2 (big nodule).

2

. Haracenter and the second

÷,

ી

# Results

9

#### *Field* varietal screening

At Bambey station the **cultivars** 83-127, Sud Cameroun, Sarakawa. 3, Sarakawa 10 and Lassa 1 had a relative effectiveness (RE), based on shoot dry weight, below 40% (Fig. 1) while at Nioro station, all cultivars had the same RE of over 40% (Fig. 2). At Bambey station, only cultivar Sarakawa 10 had and RE, based on root dry weight, below 50% at Nioro station, all cultivars had the same RE of over 50%. Relative Effectiveness, based on nodule dry weight, was much lower at Bambey than at Nioro: it was in the range of 80 to 120% at Bambey for cultivars 83-131, Ketao 3, Alheride 1, Lassa 1 and Lassa 4 and in the range of 360 to 480% at Nioro for cultivars 83-129, 83-131, Sud Cameroun, Sarakawa 2, Ketao 9, V2 and 85002. Relative Effectiveness, based on shoot nitrogen content, was over 50% for all cultivars except for Awandjelo 1 at Bambey station.

These **results** indicated that Bambey and Nioro rhizobial population established a moderately effective symbiosis with the cultivars of bambara groundnut used. Among these cultivars, **12** grew better at both the locations without **nitrogen fertilizer** (Table 2) indicating their ability to set an effective symbiosis in the two different field conditions.

Bambara Cultivars	Bambey	Nioro	
79-1	55.5 <sup>b</sup>	30.4 <sup>6</sup>	normal analytic and the
83-126	62.2 <sup>b</sup>	33.6 <sup>b</sup>	
83-129	55.8 <sup>b</sup>	39.0 <sup>ab</sup>	
83-131	56.5 <sup>b</sup>	30.2 <sup>b</sup>	
Sarakawa 1	52.8;	31.1 <sup>b</sup>	
Awanclielo 1	64.2 <sup>°</sup>	40.5 <sup>ab</sup>	
Ketao 2	63.8 <sup>b</sup>	31.7 <sup>b</sup>	
Ketao 3	66.2 <sup>ab</sup>	32.9 <sup>b</sup>	
Alheride 1	68.2 <sup>ab</sup>	43.9ª	
Lassa 3	78.5'	38.6 <sup>ab</sup>	
v2	61.5 <sup>b</sup>	41.3 <sup>ab</sup>	
85001	56.5 <sup>b</sup>	42.0"	

Table 2 Shoot dry weight (g/10 plants) of the **performed** cultivars of bambara groundnut in field at Bambey and Nioro **experimental** stations

Values followed by the **same** letter in **each column** do not **differ significantly** at the 0.05 **level** by **DUNCAN's** multiple range test (1955).

# Effectiveness of indigenous Rhizobium strains isolated from bambara groundnut nodules

Bambara groundnut cultivars sampled were moderately nodulated on both the tap and lateral roots. Many nodules appeared to be effective as indicated by interior colour. Thirty one isolates were so obtained: ( $I_1$  to  $I_{31}$ , Fig. 3). The isolates were slowgrowing and formed small (diam. < 1 mm), dry colonies on yeast mannitol agar. Four isolates ( $I_2$ ,  $I_{13}$ ,  $I_{22}$  and  $I_{30}$ ) were lodged in West Africa MIRCEN (MAO) *Rhizobium* culture collection because of their high effectiveness with cultivar 85001 which was from Senegal and had good RE values at both Bambey and Nioro experimental





and the state of the state of the

 $\mathbb{C}^{n}$ \* a

R. C. S. S. Same, I. S. S.

-

Ę

۵.

Ŧ





station. These isolates were named MAO 113, MAO 118, MAO 121 and MAO 126, respectively, and are indicated by an asterisk in Fig. 3.

#### Evaluation of host cultivar-Rhizobium strains interactions

*Rhizobium* strains used were compared for their effectiveness to nodulate the most promising cultivars of bambara groundnut selected in field varietal screening. Nodules were found in the top 5 cm of root system on all plants except controls which were all nodule free. Nodulation index and the effectiveness varied greatly with the strains (Table 3). Strains MAO 113 and MAO 118 exhibited effective nodulation on six and three cultivars respectively while strains TAL 22, TAL 169 and TAL 569 were effective on five, four and three cultivars respectively. All Rhizobium strains were effective on cultivar 83-131 exceptstrain TAL 169 which exhibited partial nodulation. All *Rhizobium* strains were partially effective on cultivar Ketao 3. Similarly, Rhizobium strains MAO 113 and MAO 118 were ineffective on cultivar V2 while strains TAL 22, TAL 169 and TAL 569 were partially effective on this cultivar. Strains MAO 113 and TAL 169 were ineffective on cultivars Awandjelo 1 and 83-129 respectively; strain MAO 118 was ineffective on cultivars Awandjelo 1, Lassa 3 and V2; strain TAL 569 was ineffective on cultivars Ketao 2, Alheride 1 and Lassa 3, Strain TAL 22 presented the widest effectiveness spectrum: it was fully effective on 42% and partially effective on 58% of all cultivars studied, followed by strain MAO 113. These results indicated that plant genotype had an important effect on nodulation of bambara gioundnut.

Although Corbin *et al.*, (1977) had proposed a scoring system to classify nodulation and Bordeleau *et al.*, (1981) had reported a method for scoring nodulation, we adopted another method. The reliability of this method was indicated by the fact that nodulation index and shoot dry weight (Table 3) showed the same trend of effectiveness in host plant-Rhizobiurn strain combinations.

#### Discussion

anda Alteration of the

+ is,

Many studies have reported an improvement of biological nitrogen fixation resulting from legume cultivar selection (Heichel, 1982; Minchin et al., 1978; Zary et al., 1978), However, selection for increased nitrogen fixation cannot be achieved by selection of the legume alone; both host genotype and rhizobial strain influence nitrogen fixation (Arrendell, et al., 1985; Hohenberg et al., 1982; Isleib et al., 1980; Zary et al., 1978). Most of the studies reported on plant genotype Rhizobium strain interactions indicated differences between cultivars of the same species in ability to fix nitrogen using introduced rhizobial strains. Because it is difficult to introduce and establish new *Rhizobium* strains into fields containing indigenous strains (Brockwell, 1980), it is important to **isolate competitive** and adapted strains, which **can** nodulate effectively most cultivars even strains are not widely distributed in the soil. Instead of introducing new Rhizobium strains, the isolated adapted Rhizobium strains may be re-introduced into the field so that they will compete against indigenous strains which are less numerous. Our results show that all Rhizobr'um strains including introduced (NifTAL) and indigenous strains nodulated all cultivars. When the hosts and rhizobial strains were arranged in an increasing order of effectiveness, one indigenous Rhizobium MAO 113 exhibited the highest effective association with the cultivars.

6



Fig. 3 Shoot dry weight of Voandzeia subterranea cultivated in Leonard jar assembly and inoculated with 31 rhizobial isolates. Treatments with a differ significantly with the control at the 0.05 level by DUNCAN's multiple range test (1955).

 7

Bambara Cultivars	Control		MAO 113 N		··· - MAC	IAO 118 TA		L 22 " TAL		169 TAL 569		
	SDW'	Nod. 1.'	SDW	Nod. 1.	SDW	Nod. 1.	SDW	Nod. I.	SDW	Nod. I.	SDW	Nod. I.
79-1	0.9ª	nn	1.4 <sup>abc</sup>	e	2.1ª	Е	2.0ª	E	1.8ª	E	1.8ª -	- <u>    e                                </u>
83-126	1.1'	nn	1.5"""	E	1.6''''	с	1.4 <sup>bc</sup>	e	1.8'	E	1.6 <sup>ab</sup>	c
83-129	1.0'	nn	1.6 <sup>ab</sup>	Е	1.5 <sup>abc</sup>	e	1.4 <sup>bc</sup>	e	1.1 <sup>bc</sup>	I	1.9"	E
83-131	1.0"	nn	1.6 <sup>ab</sup>	Е	1.8 <sup>ab</sup>	F	1.8""	Е	1.5 <sup>ab</sup>	с	1.8"	Е
Sarakawa 1	1.2"	nn	1.6 <sup>ab</sup>	Е	1.6""	е	1.8 <sup>ab</sup>	E	1.5 <sup>ab</sup>	e	1.5 <sup>ab</sup>	e
Awandjelo 1	0.9"	nn	1.1'	1	1.2""	I	1.5 <sup>abc</sup>	e	1.9ª	E	1.5 <sup>ab</sup>	e
Ketao 2	0.9"	nn	1.6""	Е	1,8 <sup>ab</sup>	с	$1.5^{abc}$	e	1.5 <sup>ab</sup>	с	1.1 <sup>b</sup>	I
Ketao 3	1.2"	nn	1.3abc	е	1.6 <sup>abc</sup>	c	1.4 <sup>bc</sup>	e	1.5 <sup>ab</sup>	e	1.6 <sup>ab</sup>	e
Alheride 1	1.2"	n n	t.3ªbc	<b>C</b> .	" 1.6 <sup>abc</sup>	e	1.8 <sup>ab</sup>	É	14 <sup>abc</sup>	e	1.1 <sup>b</sup>	i
Lassa 3	1.0*	nn	1.7"	Ē	1.1 <sup>b</sup>	1	1.9""	Ē	1.1 <sup>bc</sup>	e	1.1"	I
v2	1.0"	nn	1.5 <sup>abc</sup>	1	1.1 <sup>b</sup>	Ī	1.2°	e	0.9°	е	1.5 <sup>ab</sup>	e
85001	0.9*	nn	1.4 <sup>abc</sup>	e	2.0"	Е	1 5 <sup>ab</sup>	е	1.9'	Ē	1.8'	E

Table 3 Shoot dry weight (SDW) (g/plant) and nodulation index (Nod. I.) of 12 bambara groundnut cultivars inoculated by cowpea strains from West MIRCEN Africa (MAO) and NifTAL project

<sup>1</sup> Values followed by the same letter in each column do not differ at the 0.05 level by Duncan's multiple range test (1955). a: highly effective; ab: partially effective; b: ineffective.

<sup>2</sup> nn: non-nodulated; E: highly effective nodlation ( $\geq 4$ ); e: partially effective nodulation ( $2 \leq \text{Nod. I.} < 4$ ); I: ineffective nodulation (Nod. I. < 2).

. . .

 $\sim$ 

ł

A ....

÷.

X

Therefore, strain MAO 113 is expected to give better **nitrogen** fixation under field conditions.

م توجيع در م

The strategy to improve **nitrogen** fixation by legumes in **general** is to **first** improve genotype and then select for best *Rhizobium* strains to increase **nitrogen** fixation. Selection of host plants with good general effectiveness in nitrogen fixing ability is an other avenue (Wynne et al. 1980). The strategy most likely is to select in each field condition the most effective nitrogen fixing host plant-Rhizobium strain combination. In our study, the higher relative effectiveness obtained with cultivars at Nioro experimental station might be due to better soil conditions (higher organic matter) that are favourable to this plant (Tardieux, 1975). However, the determination of the relative effectiveness of host plant-findigenous Rhizobium strains combinations permitted us to select the best combinations in each area. Greenhouse evaluation of these combinations is the initial phase of the process of increasing **nitrogen** fixation through *Rhizobium* strains selection. The second phase was to investigate field response of bambara g.roundnut to inoculation with the selected *Rhizobium* strains. Effect of inoculation with strains MAO 113, MAO 118, TAL 22 and TAL 569 on the grain yield of 79-1 and 83-131 was obtained in two Senegalese soil (Gueye) 1987. However, longterm efforts will be required to breed for improving nitrogen fixation in bambara groundnut.

#### Acknowledgements

This research was supported by grant no. BNF-SN-2-84-21 from the U.S. **BOSTID**-National Academy of Sciences. The authors thank Alioune GNING and Omar TOURE for valuable technical assistance.

#### References

- ARRENDELL, S., WYNNE, J. C., ELKAN, G. H. and ISLEIB, T. G. 1985 Variation for nitrogen fixation progenies of a Virginia x Spanish peanut cross. 25, 865–869.
- BORDELEAU, L. M., GIROUX, M., OUELLET, R. and ANTOUN, H. 1981 Effet du soufre et de l'azote sur la fixation symbiotique d'azote chez les plantules de luzerne (Medicugo sativa L.). Canadian Journal of Plant Science 61, 639-645.
- BROCKWELL, J. 1980 Experiments with crop and pasture legumes Principles and practice. In Methodes for evaluating biological nitrogen fixation. Edited by Bergersen, F. J. pp. 417-490. Chichester, New York, Brisbane, Toronto: John Wiley and Sons.
- CALDWELL, B. E. 1966 Inheritance of a strain-specific ineffective nodulation in soybean. Crop Science 6, 427-428.
- CORBIN, E. J., BROCKWELL, J. and GAULT, R. R. 1977 Nodulation studies on chickpea (Cicer arietinum). Australian Journal of Experimental Agricultural Husbandry 17, 124-134.
- DUNCAN, D. B. 1955 Multiple range and multiple test. *Biometrics* 11, i-42.
- GUEYE, M. 1987 Effect of inoculation with different *Rhizobium* strains on grain yield of bambara groundnut (*Voandzeia subterranea* (L.) Thouars) in two senegalese soils. Paper presented at the symposium on the contribution of biological nitrogen fixation to plant production. 3-7 August, 1987. Bogor, Indonesia.
  HEICHEL, G. H. 1982 Breeding alfalfa for improved nitrogen fixation: a physiological
- **HEICHEL**, G. H. 1982 Breeding alfalta for improved nitrogen fixation: a physiological perspective. *Iowa State Journal of Research* 56, 255–280.
- **HewITT, E. J.** 1966 Sand and water culture methods **used** in the study of plant nutrition. Technical communication No. 22 (2nd ed.) 547 pp. **Commonwealth agricultural** bureaux. London.

HOHENBERG, J. S., MUNNS, D. N. and TUCKER, C. L. 1982 Rhizobium-host specificities in Phaseolus coccineus L. and Phaseolus vulgaris L. Crop Science 22, 445-459.

이 이상이 속도 잘 같아. 감사 물

ISLEIB, T. G., WYNNE, J. C., ELKAN, G. H. and SCHNEEWEIS, T. J. 1980 Quantitative genetics aspects of nitrogen fixation in peanut (Arachis hypogaea L.). Peanut Science 7, 101-105.

LEONARD, L. T. 1943 A simple assembly for use in the testing of culture of rhizobia. Journal of Bacteriology 45, 523-527.

- MINCHIN, F. R., SUMMERFIELD, R. J. and EAGLESHAM, A. R. J. 1978 Plant genotype Rhirobium strains interactions in cowpea Vigna unguiculata (L.) Walp.). Tropical Agriculture (Trinidad) 55.107-115.
- MULONGOY, K., AYANABA, A. and PULVER, E. 1980 Exploiting the diversity in the cowpearhizohia symbiosis for increased cowpea production. In Proceedings of the Global Impact of Applied Microhiology held at Lagos; Part 2. Rhizosphere biology. pp. 121-125. Academic Press. London.
- NATIONAL ACADEMY OF SCIENCES, 1979 Pulses. In Tropical Legumes: resources for the future. pp. 47-53. National Academy of Sciences, Washington D.C.
- OLSEN, S. R., C~LE, L. V., WATANABE, F. S. and DEAN, L. A. 1954 Estimation of available phosphate in soils by extraction with sodium bicarbonate. Circular of U.S. Department of Agriculture No. 939.
- SEETIN, M. W. and BARNES, D. K. 1977 Variation among alfalfa genotypes for acetylene reduction. Crop Science 17, 783-787.
- TARDIEU, M. 1957 les cultures d'appoint dans la zone du CRA Bambey. Bulletin Agronomique. Annules du Centre de Recherches Agronomiques de Bumbey au Senegal 17, 5-54.
- VINCENT, J. M. 1970 A manual for the practical study of the root-nodule bacteria. I.B.P. Handbook No. 15 Blackwell Scientific Publications. Oxford and Edinburgh. 164 pp.
- WYNNE, J. C., ELKAN, G. H., MEISNER, C. M., SCHNEEWEIS, T. J. and LIGON, J. M. 1980
- Greenhouse evaluation of strains of *Rhizobium* for peanut. *Agronomy Journal* 72, 645-649. ZARY, K. W., MILLER, Jr., J. C., WEAVER, R. W. and BARNES, L. W. 1978 Intraspecific variability for nitrogen fixation in southernpea (Vigna unguiculata (L.) Walp). Journal of American Society for Horticulture Science 103, 806-808.

#### Summarv

To improve biological nitrogen fixation in bambara groundut (Voandzeia subterranea (L.) Thouars), the relative effectiveness of 24 cultivars was studied in the field at two experimental stations selected for their different ecological conditions. Thirty one Rhizobium strains were isolated and 12 cultivars were screened during this study, Both indigenous and introduced Niftal Rhizobium strains were used during a second study on host cultivars X Rhizobium strains interactions. Nodulation index and shoot dry weight were used to assess the efficiency. The widest effectiveness spectrum was obscrved with thr indigenous strain MAO 113 and the introduced strain TAL 22.

#### Résumé

#### Fixation d'azote dans la noix de Batnbara (Voandzeia subterrana (L.) Thouars)

Dans le but d'améliorer la fixation biologique d'azote dans la noix de Bambara (Voandzeia subterranea (L.) Thouars), l'efficacité relative de 24 cultivars a été étudiée sur le terrain dans deux stations expérimentales choisies pour leurs conditions écologiques différentes. Trente-etune souches de Rhizobium ont été isolées et 12 cultivnrs ont été testées au cours de cette étude. Tant les souches indigènes de Rhizobium que celles Niftal introduites ont été utilisées au cours d'une deuxième étude sur les interractions entre les souches de Rhizobium et les cultivarshôtes. L'indice de nodulation et le poids sec de la pousse ont été utilisées pour l'établissement de l'efficience. Le spectre d'efficience le plus large a été observé avec la souche indigène MAO 113 et la souche introduite TAI. 22.

\*

18. ja 11. ja

# Resumen

Afin de mejorar la fijación biológica de nitrógeno en el cacahuete de Bambara (Voandzeia subterranea (L.) Thouars), se realizó un estudio de su efectividad relativa en campo en dos estaciones experimentales seleccionadas por sus distintas condiciones ecológicas. Durante este estudio se aislaron treintaiuna cepas de Rhizobium y se probaron doce cultivares del huésped. En un segundo ensayo se estudiaron las interacciones huésped xcepa de Rhizobium para lo cual se utilizaron cepas de Rhizobium Niftal tanto indígenas como introducidas. La eficiencia se valoró determinando el indice de nodulación y el peso seco de la parte aerea. El espectro mas amplio de eficacia se observó con la cepa indígena MAO 113 y la introducida TAL 22.